

- Gibson, C. E., Smith, R. V. & Stewart, D. A. 1988. A long term study of the phosphorus cycle in Lough Neagh, Northern Ireland. *Int. Revue Ges. Hydrobiol.* **73**(3): 249–257.
- Langton, P. H. 1984. *A key to pupal exuviae of British Chironomidae*. 324 pp. Privately published.
- Langton, P. H. 1991. *A key to pupal exuviae of West Palaearctic Chironomidae*. 386 pp. Privately published.
- Langton, P. H. 1993. Species additional to those included in Pinder 1978. *The Edwardian, British Chironomists' Newsletter*, 4 pp. (not numbered).
- Murray, J. & Pullar, L. 1977. *Bathymetrical survey of Scottish fresh-water lochs*. Arno Press.
- OECD 1982. *Eutrophication of waters. Monitoring, assessment and control*. 154 pp. OECD Paris.
- Reiss, F. 1968. Ökologische und systematische Untersuchungen an Chironomiden (Diptera) des Bodensees. Ein Beitrag zur lakustrischen Chironomidenfauna des nördlichen Alpenvorlandes. *Arch. Hydrobiol.* **64**: 176–323.
- Pagast, F. 1947. Systematik und Verbreitung der um die Gattung *Diamesa* gruppierten Chironomiden. *Arch. Hydrobiol.* **41**: 435–596.
- Sæther, O. A. 1973. Taxonomy and ecology of three new species of *Monodiamesa* Kieffer, with keys to nearctic and palaearctic species of the genus (Diptera: Chironomidae). *J. Fish. Res. Bd. Can.* **30**: 665–679.
- Sæther, O. A. 1977. Female genitalia in Chironomidae and other Nematocera: morphology, phylogenies, keys. *Bull. Fish. Res. Bd. Can.* **197**: 209 pp.
- Sæther, O. A. 1983. 8. The larvae of Prodiamesinae (Diptera: Chironomidae) of the Holarctic region. Keys and diagnoses. *Ent. Scand. Suppl.* **19**: 141–147.
- Stevens, R. J. & Gibson, C. E. 1977. Sediment release of phosphorus in Lough Neagh, Northern Ireland. pp. 343–347. In: Golterman, H. L. (Ed.) *Interactions between sediments and freshwater*. D. W. Junk B. V. Publishers, The Hague.

SHORT COMMUNICATION

Dryinus pumilio Puton (Hemiptera: Lygaeidae) in Surrey.—A single example of this rare bug was swept from rideside weedy grassland in Preserve Cope immediately to the east of Polesden Lacey (TQ138522), 10.ix.1996. This wood is probably of plantation origin, having at present mature standard oaks of very uniform age plus a mixture of exotic tree species. The canopy is very sparse due to severe storm damage in 1987 and subsequent heavy suppression of any regrowth by rabbit grazing. The wood lies on the clay-with-flints cap of the North Downs and the soils are neutral. The grassland includes frequent *Hypericum perforatum*, a feature of other sites mentioned by Kirby (1992). This is a nationally scarce bug with a thin scattering of reports across lowland southern Britain from Kent to Monmouthshire, and equally scarce throughout its range—it appears to be confined to western Europe. So far as I am aware this is the first record from Surrey.—K. N. A. ALEXANDER, National Trust, 33 Sheep Street, Cirencester, Glos. GL7 1RQ.

REFERENCES

- Kirby, P. 1992. *A review of the scarce and threatened Hemiptera of Great Britain*. UK Nature Conservation No. 2. Joint Nature Conservation Committee, Peterborough.